Index to Volume 9: Subject

A

Acoustics
Acoustic approaches to remote species identification: a review (Horne), 356

Adriatic Sea
Climate change and Adriatic ichthyofauna (Dulcic *et al.*), 187

Aggregation
On the use of Lloyd's index of patchiness (Bez), 372

Amundsen Sea
Distribution of minke whales in the Bellinghausen and Amundsen Seas (60°W–120°W), with special reference to environmental/physiographic variables (Kasamatsu *et al.*), 214

Anchovy
Alternating dominance of postlarval sardine and anchovy caught by coastal fishery in relation to the Kuroshio meander in the Enshu-nada Sea (Nakata *et al.*), 248

Arguin bank
The importance of retention process in upwelling areas for recruitment of *Octopus vulgaris*: the example of the Arguin bank (Mauritania) (Faure *et al.*), 343

Atlantic bluefin tuna
Schooling and migration of large pelagic fishes relative to environmental cues (Humston *et al.*), 136

Atlantic herring
Observations on larval fish transport and retention on the Scotian Shelf in relation to geostrophic circulation (Reiss *et al.*), 195

B

Backscatter amplitude
Acoustic approaches to remote species identification: a review (Horne), 356

Bakun index
Temporal and spatial responses of British Columbia steelhead (*Oncorhynchus mykiss*) populations to ocean climate shifts (Welch *et al.*), 17

Barents Sea
Predicting the temperature of the Barents Sea (Ottersen *et al.*), 121

Behaviour
Behaviour of immature yellowtails (*Seriola quinqueradiata*) observed by electronic data-recording tags (Kasai *et al.*), 259
A dynamic optimisation model for the behaviour of tunas at ocean fronts (Kirby *et al.*), 328

Bellinghausen Sea
Distribution of minke whales in the Bellinghausen and Amundsen Seas (60°W–120°W), with special reference to environmental/physiographic variables (Kasamatsu *et al.*), 214

Bering Sea
A synthesis of biological and physical processes affecting the feeding environment of larval walleye pollock (*Theragra chalcogramma*) in the eastern Bering Sea (Napp *et al.*), 147
Diurnal variation in thermal environment experienced by salmonids in the North Pacific as indicated by data storage tags (Walker *et al.*), 171
Pacific herring, *Clupea pallasi*, recruitment in the Bering Sea and north-east Pacific Ocean, I: relationships among different populations (Williams *et al.*), 285
Pacific herring, *Clupea pallasi*, recruitment in the Bering Sea and north-east Pacific Ocean, II: relationships to environmental variables and implications for forecasting (Williams *et al.*), 300

Bioenergetics
Computer simulations of the effects of the Sitka eddy on the migration of sockeye salmon returning to British Columbia (Healey *et al.*), 271

C

California Current
Coastal sea surface temperature and coho salmon production off the north west United States (Cole), 1

Caretta caretta
Turtles on the edge: movement of loggerhead turtles (*Caretta caretta*) along oceanic fronts, spanning longline fishing grounds in the central North Pacific, 1997–1998 (Polovina *et al.*), 71

Catchability
Environmental influences on the availability of smooth pink shrimp, *Pandalus jordani*, to commercial fishing gear off Vancouver Island, Canada (Perry *et al.*), 50

Central North Pacific
Turtles on the edge: movement of loggerhead turtles (*Caretta caretta*) along oceanic fronts, spanning longline fishing grounds in the central North Pacific, 1997–1998 (Polovina *et al.*), 71

Circulation
A synthesis of biological and physical processes affecting the feeding environment of larval walleye pollock (*Theragra chalcogramma*) in the eastern Bering Sea (Napp *et al.*), 147
Observations on larval fish transport and retention on the Scotian Shelf in relation to geostrophic circulation (Reiss *et al.*), 195

Climate change
Climate change and Adriatic ichthyofauna (Dulcic *et al.*), 187

Climate effects
Decadel-scale patterns in the relative year class success of sablefish (*Anoplopoma fimbria*) (King *et al.*), 62

Climatic warming
Distribution of minke whales in the Bellingshausen and Amundsen Seas (60°W–120°W), with special reference to environmental/physiographic variables (Kasamatsu *et al.*), 214

Clupea harengus
Observations on larval fish transport and retention on the Scotian Shelf in relation to geostrophic circulation (Reiss *et al.*), 195

Coastal retention
The importance of retention process in upwelling areas for recruitment of *Octopus vulgaris*: the example of the Arguin bank (Mauritania) (Faure *et al.*), 343

Coastal upwelling
The importance of retention process in upwelling areas for recruitment of *Octopus vulgaris*: the example of the Arguin bank (Mauritania) (Faure *et al.*), 343

Cod
Patterns in the spawning of cod (*Gadus morhua* L.), sole (*Solea solea* L.) and plaice (*Pleuronectes platessa* L.) in the Irish Sea as determined by generalized additive modelling (Fox *et al.*), 33
Observations on larval fish transport and retention on the Scotian Shelf in relation to geostrophic circulation (Reiss *et al.*), 195
Coho
Temporal and spatial responses of British Columbia steelhead (*Oncorhynchus mykiss*) populations to ocean climate shifts (Welch *et al.*), 17
Coho salmon
Coastal sea surface temperature and coho salmon production off the north west United States (Cole), 1
Counts
On the use of Lloyd's index of patchiness (Bez), 372
CPUE
Environmental influences on the availability of smooth pink shrimp, *Pandalus jordani*, to commercial fishing gear off Vancouver Island, Canada (Perry *et al.*), 50
Crowding
On the use of Lloyd's index of patchiness (Bez), 372

D
Data recording tag
Behaviour of immature yellowtails (*Seriola quinqueradiata*) observed by electronic data-recording tags (Kasai *et al.*), 259
Data storage tags
Diurnal variation in thermal environment experienced by salmonids in the North Pacific as indicated by data storage tags (Walker *et al.*), 171
Decadal change
Ecological variations and El Niño effects off the southern coast of the Korean Peninsula during the last three decades (Kim *et al.*), 239
Densities
On the use of Lloyd's index of patchiness (Bez), 373
Diagnostic calculation
Observations on larval fish transport and retention on the Scotian Shelf in relation to geostrophic circulation (Reiss *et al.*), 195
Distribution
Distribution of minke whales in the Bellingshausen and Amundsen Seas (60°W–120°W), with special reference to environmental/physiographic variables (Kasamatsu *et al.*), 214
Dominance alternation
Alternating dominance of postlarval sardine and anchovy caught by coastal fishery in relation to the Kuroshio meander in the Enshu-nada Sea (Nakata *et al.*), 248
Driftnet fishing
Interannual variability in abundance of the neon flying squid, *Ommastrephes bartramii*, in the North Pacific Ocean during 1979-1998: impact of driftnet fishing and oceanographic conditions (Yatsu *et al.*), 163

E
El Nino effect
Ecological variations and El Niño effects off the southern coast of the Korean Peninsula during the last three decades (Kim *et al.*), 239
Engraulis japonica
Alternating dominance of postlarval sardine and anchovy caught by coastal fishery in relation to the Kuroshio meander in the Enshu-nada Sea (Nakata *et al.*), 248
Enshu-nada Sea
Alternating dominance of postlarval sardine and anchovy caught by coastal fishery in relation to the Kuroshio meander in the Enshu-nada Sea (Nakata *et al.*), 248
Environmental variable
Pacific herring, *Clupea pallasi*, recruitment in the Bering Sea and north-east Pacific Ocean, II: relationships to environmental variables and implications for forecasting (Williams *et al.*), 300

F
Fisheries oceanography
A synthesis of biological and physical processes affecting the feeding environment of larval walleye pollock (*Theragra chalcogramma*) in the eastern Bering Sea (Napp *et al.*), 147
Fishing gear
Environmental influences on the availability of smooth pink shrimp, *Pandalus jordani*, to commercial fishing gear off Vancouver Island, Canada (Perry *et al.*), 50
Forecasting
Pacific herring, *Clupea pallasi*, recruitment in the Bering Sea and north-east Pacific Ocean, II: relationships to environmental variables and implications for forecasting (Williams *et al.*), 300
Fronts
A dynamic optimisation model for the behaviour of tunas at ocean fronts (Kirby *et al.*), 328

G
Gadus morhua
Patterns in the spawning of cod (*Gadus morhua* L.), sole (*Solea solea* L.) and plaice (*Pleuronectes platessa* L.) in the Irish Sea as determined by generalized additive modelling (Fox *et al.*), 33
Observations on larval fish transport and retention on the Scotian Shelf in relation to geostrophic circulation (Reiss *et al.*), 195
Generalized additive modelling
Patterns in the spawning of cod (*Gadus morhua* L.), sole (*Solea solea* L.) and plaice (*Pleuronectes platessa* L.) in the Irish Sea as determined by generalized additive modelling (Fox *et al.*), 33
Growth
Interannual variability in abundance of the neon flying squid, *Ommastrephes bartramii*, in the North Pacific Ocean during 1979-1998: impact of driftnet fishing and oceanographic conditions (Yatsu *et al.*), 163
Gulf of Maine
Modelling the transport of lobster (*Homarus americanus*) larvae and postlarvae in the Gulf of Maine (Incze *et al.*), 99

H
Hake
Observations on larval fish transport and retention on the Scotian Shelf in relation to geostrophic circulation (Reiss *et al.*), 195
Homarus americanus
Modelling the transport of lobster (*Homarus americanus*) larvae and postlarvae in the Gulf of Maine (Incze *et al.*), 99

I
Ice
Distribution of minke whales in the Bellinghausen and Amundsen Seas (60°W–120°W), with special reference to environmental/physiographic variables (Kasamatsu *et al.*), 214
Ichthyofauna
Climate change and Adriatic ichthyofauna (Dulcic *et al.*), 187
Individual-based modelling
Computer simulations of the effects of the Sitka eddy on the migration of sockeye salmon returning to British Columbia (Healey *et al.*), 271
Irish Sea
Patterns in the spawning of cod (*Gadus morhua* L.), sole (*Solea solea* L.) and plaice (*Pleuronectes platessa* L.) in the Irish Sea as determined by generalized additive modelling (Fox *et al.*), 33

K
Kinesis
Schooling and migration of large pelagic fishes relative to environmental cues (Humston *et al.*), 136
Kuroshio meander
Alternating dominance of postlarval sardine and anchovy caught by coastal fishery in relation to the Kuroshio meander in the Enshu-nada Sea (Nakata *et al.*), 248

L
Larvae
Modelling the transport of lobster (*Homarus americanus*) larvae and postlarvae in the Gulf of Maine (Incze *et al.*), 99
Larval dispersal
Spatial variability in red sea urchin (*Strongylocentrotus franciscanus*) recruitment in northern California (Morgan *et al.*), 83
Larval feeding
A synthesis of biological and physical processes affecting the feeding environment of larval walleye pollock (*Theragra chalcogramma*) in the eastern Bering Sea (Napp *et al.*), 147
Larval fish survival
A synthesis of biological and physical processes affecting the feeding environment of larval walleye pollock (*Theragra chalcogramma*) in the eastern Bering Sea (Napp *et al.*), 147
Lobster
Modelling the transport of lobster (*Homarus americanus*) larvae and postlarvae in the Gulf of Maine (Incze *et al.*), 99
Loggerhead turtles
Turtles on the edge: movement of loggerhead turtles (*Caretta caretta*) along oceanic fronts, spanning longline fishing grounds in the central North Pacific, 1997–1998 (Polovina *et al.*), 71
Longline fishery
Turtles on the edge: movement of loggerhead turtles (*Caretta caretta*) along oceanic fronts, spanning longline fishing grounds in the central North Pacific, 1997–1998 (Polovina *et al.*), 71

M
Marine banks
Observations on larval fish transport and retention on the Scotian Shelf in relation to geostrophic circulation (Reiss *et al.*), 195
Marine ecosystem
Ecological variations and El Niño effects off the southern coast of the Korean Peninsula during the last three decades (Kim *et al.*), 239
Marine survival
Coastal sea surface temperature and coho salmon production off the north west United States (Cole), 1
Merluccius billinearis
Observations on larval fish transport and retention on the Scotian Shelf in relation to geostrophic circulation (Reiss *et al.*), 195
Meroplankton
Spatial variability in red sea urchin (*Strongylocentrotus franciscanus*) recruitment in northern California (Morgan *et al.*), 83
Mesoscale eddies
Computer simulations of the effects of the Sitka eddy on the migration of sockeye salmon returning to British Columbia (Healey *et al.*), 271
Migration
Schooling and migration of large pelagic fishes relative to environmental cues (Humston *et al.*), 136
Minke whale
Distribution of minke whales in the Bellinghausen and Amundsen Seas (60°W–120°W), with special reference to environmental/physiographic variables (Kasamatsu *et al.*), 214
Multivariate cluster analysis
Pacific herring, *Clupea pallasi*, recruitment in the Bering Sea and north-east Pacific Ocean, I: relationships among different populations (Wililams *et al.*), 285

N
Neon flying squid
Interannual variability in abundance of the neon flying squid, *Ommastrephes bartramii*, in the North Pacific Ocean during 1979-1998: impact of driftnet fishing and oceanographic conditions (Yatsu *et al.*), 163
North Pacific
Decadel-scale patterns in the relative year class success of sablefish (*Anoplopoma fimbria*) (King *et al.*), 62
Interannual variability in abundance of the neon flying squid, *Ommastrephes bartramii*, in the North Pacific Ocean during 1979-1998: impact of driftnet fishing and oceanographic conditions (Yatsu *et al.*), 163
Diurnal variation in thermal environment experienced by salmonids in the North Pacific as indicated by data storage tags (Walker *et al.*), 171
Mid-latitude wind stress: the energy source for climatic shifts in the North Pacific Ocean (Parrish *et al.*), 224
North-east Pacific Ocean
Pacific herring, *Clupea pallasi*, recruitment in the Bering Sea and north-east Pacific Ocean, I: relationships among different populations (Williams *et al.*), 285
Pacific herring, *Clupea pallasi*, recruitment in the Bering Sea and north-east Pacific Ocean, II: relationships to environmental variables and implications for forecasting (Williams *et al.*), 300
Northern California
Spatial variability in red sea urchin (*Strongylocentrotus franciscanus*) recruitment in northern California (Morgan *et al.*), 83

Numerical models
Modelling the transport of lobster (*Homarus americanus*) larvae and postlarvae in the Gulf of Maine (Incze *et al.*), 99
Nursery habitat
Relative contributions from exposed inshore and estuarine nursery grounds to the recruitment of stone flounder, *Platichythys bicoloratus*, estimated using otolith Sr:Ca ratios (Yamashita *et al.*), 316

O
Ocean climate
Temporal and spatial responses of British Columbia steelhead (*Oncorhynchus mykiss*) populations to ocean climate shifts (Welch *et al.*), 17
Oceanographic variability
Distribution of minke whales in the Bellinghausen and Amundsen Seas (60°W–120°W), with special reference to environmental/physiographic variables (Kasamatsu *et al.*), 214
Octopus vulgaris
The importance of retention process in upwelling areas for recruitment of *Octopus vulgaris*: the example of the Arguin bank (Mauritania) (Faure *et al.*), 345
Operational oceanography
Observations on larval fish transport and retention on the Scotian Shelf in relation to geostrophic circulation (Reiss *et al.*), 195
Optimal interpolation
Observations on larval fish transport and retention on the Scotian Shelf in relation to geostrophic circulation (Reiss *et al.*), 195
Otolith
Relative contributions from exposed inshore and estuarine nursery grounds to the recruitment of stone flounder, *Platichythys bicoloratus*, estimated using otolith Sr:Ca ratios (Yamashita *et al.*), 316

P
Pacific herring
Pacific herring, *Clupea pallasi*, recruitment in the Bering Sea and north-east Pacific Ocean, I: relationships among different populations (Williams *et al.*), 285
Pacific herring, *Clupea pallasi*, recruitment in the Bering Sea and north-east Pacific Ocean, II: relationships to environmental variables and implications for forecasting (Williams *et al.*), 300
Pacific salmon
Diurnal variation in thermal environment experienced by salmonids in the North Pacific as indicated by data storage tags (Walker *et al.*), 171
Physiology
A dynamic optimisation model for the behaviour of tunas at ocean fronts (Kirby *et al.*), 329
Plaice
Patterns in the spawning of cod (*Gadus morhua* L.), sole (*Solea solea* L.) and plaice (*Pleuronectes platessa* L.) in the Irish Sea as determined by generalized additive modelling (Fox *et al.*), 33
Platichthys biocoloratus
Relative contributions from exposed inshore and estuarine nursery grounds to the recruitment of stone flounder, *Platichythys bicoloratus*, estimated using otolith Sr:Ca ratios (Yamashita *et al.*), 316
Pleuronectes platessa
Patterns in the spawning of cod (*Gadus morhua* L.), sole (*Solea solea* L.) and plaice (*Pleuronectes platessa* L.) in the Irish Sea as determined by generalized additive modelling (Fox *et al.*), 33
Postlarvae
Modelling the transport of lobster (*Homarus americanus*) larvae and postlarvae in the Gulf of Maine (Incze *et al.*), 99
Prediction
Predicting the temperature of the Barents Sea (Otterson), 121

R
Recruitment
Pacific herring, *Clupea pallasi*, recruitment in the Bering Sea and north-east Pacific Ocean, I: relationships among different populations (Williams *et al.*), 285
Pacific herring, *Clupea pallasi*, recruitment in the Bering Sea and north-east Pacific Ocean, II: relationships to environmental variables and implications for forecasting (Williams *et al.*), 300
Relative contributions from exposed inshore and estuarine nursery grounds to the recruitment of stone flounder, *Platichythys bicoloratus*, estimated using otolith Sr:Ca ratios (Yamashita *et al.*), 316
The importance of retention process in upwelling areas for recruitment of *Octopus vulgaris*: the example of the Arguin bank (Mauritania) (Faure *et al.*), 343
Recruitment varia
Spatial variability in red sea urchin (*Strongylocentrotus franciscanus*) recruitment in northern California (Morgan *et al.*), 83
Red sea urchin
Spatial variability in red sea urchin (*Strongylocentrotus franciscanus*) recruitment in northern California (Morgan *et al.*), 83
Regime shift
Temporal and spatial responses of British Columbia steelhead (*Oncorhynchus mykiss*) populations to ocean climate shifts (Welch *et al.*), 17
Mid-latitude wind stress: the energy source for climatic shifts in the North Pacific Ocean (Parrish *et al.*), 224
Regimes
Decadel-scale patterns in the relative year class success of sablefish (*Anoplopoma fimbria*) (King *et al.*), 62
Remote sensing
Coastal sea surface temperature and coho salmon production off the north west United States (Cole), 1
The importance of retention process in upwelling areas for recruitment of *Octopus vulgaris*: the example of the Arguin bank (Mauritania) (Faure *et al.*), 343
Retention
Observations on larval fish transport and retention on the Scotian Shelf in relation to geostrophic circulation (Reiss *et al.*), 195

S
Salmon
Temporal and spatial responses of British Columbia steelhead (*Oncorhynchus mykiss*) populations to ocean climate shifts (Welch *et al.*), 17
Sablefish
Decadel-scale patterns in the relative year class success of sablefish (*Anoplopoma fimbria*) (King *et al.*), 62

Sardine
Alternating dominance of postlarval sardine and anchovy caught by coastal fishery in relation to the Kuroshio meander in the Enshu-nada Sea
(Nakata *et al.*), 248
Sardinops melanostictus
Alternating dominance of postlarval sardine and anchovy caught by coastal fishery in relation to the Kuroshio meander in the Enshu-nada Sea
(Nakata *et al.*), 248
Satellite remote sensing
Turtles on the edge: movement of loggerhead turtles (*Caretta caretta*) along oceanic fronts, spanning longline fishing grounds in the central North Pacific, 1997–1998
(Polovina *et al.*), 71
Scotian Shelf
Observations on larval fish transport and retention on the Scotian Shelf in relation to geostrophic circulation
(Reiss *et al.*), 195
Sea breeze
Modelling the transport of lobster (*Homarus americanus*) larvae and postlarvae in the Gulf of Maine
(Incze *et al.*), 99
Sea ice
Distribution of minke whales in the Bellinghausen and Amundsen Seas (60°W–120°W), with special reference to environmental/physiographic variables
(Kasamatsu *et al.*), 214
Sea temperature
Distribution of minke whales in the Bellinghausen and Amundsen Seas (60°W–120°W), with special reference to environmental/physiographic variables
(Kasamatsu *et al.*), 214
Sea surface temperature
Coastal sea surface temperature and coho salmon production off the north west United States
(Cole), 1
Interannual variability in abundance of the neon flying squid, *Ommastrephes bartramii*, in the North Pacific Ocean during 1979-1998: impact of driftnet fishing and oceanographic conditions
(Yatsu *et al.*), 163
Shrimp
Environmental influences on the availability of smooth pink shrimp, *Pandalus jordani*, to commercial fishing gear off Vancouver Island, Canada
(Perry *et al.*), 50
Sitka eddy
Computer simulations of the effects of the Sitka eddy on the migration of sockeye salmon returning to British Columbia
(Healey *et al.*), 271
Sockeye salmon migration
Computer simulations of the effects of the Sitka eddy on the migration of sockeye salmon returning to British Columbia
(Healey *et al.*), 271
SOI
Ecological variations and El Niño effects off the southern coast of the Korean Peninsula during the last three decades
(Kim *et al.*), 239
Sole
Patterns in the spawning of cod (*Gadus morhua* L.), sole (*Solea solea* L.) and plaice (*Pleuronectes platessa* L.) in the Irish Sea as determined by generalized additive modelling
(Fox *et al.*), 33
Solea solea
Patterns in the spawning of cod (*Gadus morhua* L.), sole (*Solea solea* L.) and plaice (*Pleuronectes platessa* L.) in the Irish Sea as determined by generalized additive modelling
(Fox *et al.*), 33
South Sea of Korea
Ecological variations and El Niño effects off the southern coast of the Korean Peninsula during the last three decades
(Kim *et al.*), 239
Spawning
Patterns in the spawning of cod (*Gadus morhua* L.), sole (*Solea solea* L.) and plaice (*Pleuronectes platessa* L.) in the Irish Sea as determined by generalized additive modelling
(Fox *et al.*), 33
Spatial model
Schooling and migration of large pelagic fishes relative to environmental cues
(Humston *et al.*), 136
Species identification
Acoustic approaches to remote species identification: a review
(Horne), 356
Spine canopy
Spatial variability in red sea urchin (*Strongylocentrotus franciscanus*) recruitment in northern California
(Morgan *et al.*), 83
SST
Mid-latitude wind stress: the energy source for climatic shifts in the North Pacific Ocean
(Parrish *et al.*), 224
Statistical methods
Predicting the temperature of the Barents Sea
(Ottersen *et al.*), 121
Steelhead
Temporal and spatial responses of British Columbia steelhead (*Oncorhynchus mykiss*) populations to ocean climate shifts
(Welsh *et al.*), 17
Steelhead trout
Diurnal variation in thermal environment experienced by salmonids in the North Pacific as indicated by data storage tags
(Walker *et al.*), 171
Stochastic dynamic programming
A dynamic optimisation model for the behaviour of tunas at ocean fronts
(Kirby *et al.*), 328
Stock abundance
Interannual variability in abundance of the neon flying squid, *Ommastrephes bartramii*, in the North Pacific Ocean during 1979-1998: impact of driftnet fishing and oceanographic conditions
(Yatsu *et al.*), 163
Stock assessment
Environmental influences on the availability of smooth pink shrimp, *Pandalus jordani*, to commercial fishing gear off Vancouver Island, Canada
(Perry *et al.*), 50
Stone flounder
Relative contributions from exposed inshore and estuarine nursery grounds to the recruitment of stone flounder, *Platichthys bicoloratus*, estimated using otolith Sr:Ca ratios
(Yamashita *et al.*), 316
Strongylocentrotus franciscanus
Spatial variability in red sea urchin (*Strongylocentrotus franciscanus*) recruitment in northern California
(Morgan *et al.*), 83
Strontium:calcium ratios
Schooling and migration of large pelagic fishes relative to environmental cues
(Humston *et al.*), 136
Subtropical front
Turtles on the edge: movement of loggerhead turtles (*Caretta caretta*) along oceanic fronts, spanning

longline fishing grounds in the central North Pacific, 1997–1998 (Polovina *et al.*), 71
Sustainable fisheries
Mid-latitude wind stress: the energy source for climatic shifts in the North Pacific Ocean (Parrish *et al.*), 224

T
Tagging effects
Diurnal variation in thermal environment experienced by salmonids in the North Pacific as indicated by data storage tags (Walker *et al.*), 171
Target strength
Acoustic approaches to remote species identification: a review (Horne), 356
Temperature
Predicting the temperature of the Barents Sea (Ottersen *et al.*), 121
Thermal habitat
Diurnal variation in thermal environment experienced by salmonids in the North Pacific as indicated by data storage tags (Walker *et al.*), 171
Tide
Environmental influences on the availability of smooth pink shrimp, *Pandalus jordani*, to commercial fishing gear off Vancouver Island, Canada (Perry *et al.*), 50
Time series
Predicting the temperature of the Barents Sea (Ottersen *et al.*), 121
Transport
Modelling the transport of lobster (*Homarus americanus*) larvae and postlarvae in the Gulf of Maine (Incze *et al.*), 99
Observations on larval fish transport and retention on the Scotian Shelf in relation to geostrophic circulation (Reiss *et al.*), 195
Tunas
A dynamic optimisation model for the behaviour of tunas at ocean fronts (Kirby *et al.*), 328

U
Upwelling
Coastal sea surface temperature and coho salmon production off the north west United States (Cole), 1
Temporal and spatial responses of British Columbia steelhead (*Oncorhynchus mykiss*) populations to ocean climate shifts (Welsh *et al.*), 17
Upwelling relaxation
Spatial variability in red sea urchin (*Strongylocentrotus franciscanus*) recruitment in northern California (Morgan *et al.*), 83

V
Variational algorithm
Observations on larval fish transport and retention on the Scotian Shelf in relation to geostrophic circulation (Reiss *et al.*), 195

W
Weight-at-age
Pacific herring, *Clupea pallasi*, recruitment in the Bering Sea and north-east Pacific Ocean, I: relationships among different populations (Wililams *et al.*), 285
Wind
Environmental influences on the availability of smooth pink shrimp, *Pandalus jordani*, to commercial fishing gear off Vancouver Island, Canada (Perry *et al.*), 50
Wind stress
Mid-latitude wind stress: the energy source for climatic shifts in the North Pacific Ocean (Parrish *et al.*), 224

Y
Year class success
Decadel-scale patterns in the relative year class success of sablefish (*Anoplopoma fimbria*) (King *et al.*), 62
Yellowtail
Behaviour of immature yellowtails (*Seriola quinqueradiata*) observed by electronic data-recording tags (Kasai *et al.*), 259

Z
Zeroes
On the use of Lloyd's index of patchiness (Bez), 373

Index to Volume 9: Author

Ådlandsvik, Bjørn, 121
Anma, Gen, 171
Ault, Jerald S., 136
Aydin, Kerim Y., 171

Balazs, George H., 71
Beamish, R.J., 114
Beamish, R.J., 62
Bez, Nicolas, 372
Boehlert, George W., 171
Botsford, Louis W., 83
Boutillier, James A., 50

Carlson, Richard H., 171
Cole, James, 1
Cury, Philippe, 343

Davis, Nancy D., 171
Demarcq, Herve, 343
Deyoung, B., 195
Dickey-Collas, M., 33
Diehl, Jennifer M., 83
Dulcic, J., 187

Ensor, Paul, 214
Eveson, J.P., 17

Faure, Valerie, 343
Fiksen, Oyvind, 328
Foreman, Michael G.G., 50
Fox, C.J., 33

Friedland, Kevin D., 171
Funakoshi, Shigeo, 248

Grbec, B., 187

Hart, Paul J.B., 328
Healey, M.C., 271
Hinch, S.G., 271
Horne, John K., 356
Huato, L., 271
Humston, Robert, 136

Incze, Lewis S., 99
Inejih, Cheikh Abdellahi, 343
Ishida, Yukimasa, 163

Joyce, Gerald G., 214
Jr., J.D., 147

Kamei, Yoshihiko, 163
Kang, Sukyung, 239
Kasai, Akihide, 259
Kasamatsu, Fujio, 214
Kendall, A.W., 147
Kim, Suam, 239
Kimura, Noritugu, 214
King, J., 114
King, J.R., 62
Kirby, David S., 328
Kobayashi, Donald R., 71
Kuno, Masahiro, 282

Leblond, P.H., 271
Leong, Harald, 121
Livingston, Patricia, 192
Lundquist, Carolyn J., 83
Lutcavage, Molly, 136

Mcfarlane, G.A., 114
Mcfarlane, G.A., 62
Meguro, Toshimi, 163
Mendelssohn, R., 224
Mitsunaga, Yasushi, 259
Morgan, Lance E., 83
Mori, Junta, 163
Myers, Katherine W., 171

Nagasawa, Kazuya, 163
Naimie, Christopher E., 99
Nakamura, Motohiko, 248
Nakata, Hideaki, 248
Napp, J.M., 147
Nash, R.D.M., 33
Noakes, D.J., 114

O'brien, C.M., 33
Olson, Donald B., 136
Otake, Tsuguo, 316
Ottersen, Geir, 121

Panteleev, G., 195
Parker, Denise M., 71
Parrish, R.H., 224
Perry, R.Ian, 50
Pinnix, W., 114
Polovina, Jeffrey J., 71

Quinn, II, Terrance J., 285
Quinn, II, Terrance J., 300

Reiss, C.S., 195

Sakamoto, Wataru, 259
Sakurai, Yasunori, 163
Schumacher, J.D., 147
Schwing, F.B., 224
Seki, Michael P., 71
Sheng, J., 195
Smith, B.D., 17
Sweeting, R., 114

Taggart, C.T., 195
Thomson, K.A., 271

Ueno, Yasuhiro, 171
Urawa, Shigehiko, 171

Walker, Robert V., 171
Walters, C.J., 271
Ward, B.R., 17
Watanabe, Tomowo, 163
Welch, D.W., 17
Williams, Erik H., 285
Williams, Erik H., 300
Wing, Stephen R., 83

Yamada, Hideaki, 316
Yamada, Hirokatsu, 282
Yamakawa, Takashi, 282
Yamamoto, Shotaro, 259
Yamashita, Yoh, 316
Yatsu, Akihiko, 163

Contents

Volume 9 Number 1 March 2000

ORIGINAL ARTICLES

1 Coastal sea surface temperature and coho salmon production off the north west United States
James Cole

17 Temporal and spatial responses of British Columbia steelhead (*Oncorhynchus mykiss*) populations to ocean climate shifts
D. W. Welch, B. R. Ward, B. D. Smith and J. P. Eveson

33 Patterns in the spawning of cod (*Gadus morhua* L.), sole (*Solea solea* L.) and plaice (*Pleuronectes platessa* L.) in the Irish Sea as determined by generalized additive modelling
C. J. Fox, C. M. O'Brien, M. Dickey-Collas and R. D. M. Nash

50 Environmental influences on the availability of smooth pink shrimp, *Pandalus jordani*, to commercial fishing gear off Vancouver Island, Canada
R. Ian Perry, James A. Boutillier and Michael G. G. Foreman

62 Decadel-scale patterns in the relative year class success of sablefish (*Anoplopoma fimbria*)
J. R. King, G. A. McFarlane and R. J. Beamish

71 Turtles on the edge: movement of loggerhead turtles (*Caretta caretta*) along oceanic fronts, spanning longline fishing grounds in the central North Pacific, 1997–1998
Jeffrey J. Polovina, Donald R. Kobayashi, Denise M. Parker, Michael P. Seki and George H. Balazs

83 Spatial variability in red sea urchin (*Strongylocentrotus franciscanus*) recruitment in northern California
Lance E. Morgan, Stephen R. Wing, Louis W. Botsford, Carolyn J. Lundquist and Jennifer M. Diehl

99 Modelling the transport of lobster (*Homarus americanus*) larvae and postlarvae in the Gulf of Maine
Lewis S. Incze and Christopher E. Naimie

114 Trends in coho marine survival in relation to the regime concept
R. J. Beamish, D. J. Noakes, G. A. McFarlane, W. Pinnix, R. Sweeting and J. King

120 ERRATUM

Contents

Volume 9 Number 2 June 2000

ORIGINAL ARTICLES

121 Predicting the temperature of the Barents Sea
Geir Ottersen, Bjørn Ådlandsvik and Harald Loeng

136 Schooling and migration of large pelagic fishes relative to environmental cues
Robert Hunston, Jerald S. Ault, Molly Lutcavage and Donald B. Olson

147 A synthesis of biological and physical processes affecting the feeding environment of larval walleye pollock (*Theragra chalcogramma*) in the eastern Bering Sea
J. M. Napp, A. W. Kendall, Jr. and J. D. Schumacher

163 Interannual variability in abundance of the neon flying squid, *Ommastrephes bartramii*, in the North Pacific Ocean during 1979–1998: impact of driftnet fishing and oceanographic conditions
Akihiko Yatsu, Tomowo Watanabe, Junta Mori, Kazuya Nagasawa, Yukimasa Ishida, Toshimi Meguro, Yoshihiko Kamei and Yasunori Sakurai

171 Diurnal variation in thermal environment experienced by salmonids in the North Pacific as indicated by data storage tags
Robert V. Walker, Katherine W. Myers, Nancy D. Davis, Kerim Y. Aydin, Kevin D. Friedland, H. Richard Carlson, George W. Boehlert, Shigehiko Urawa, Yasuhiro Ueno and Gen Anma

SHORT COMMUNICATION

187 Climate change and Adriatic ichthyofauna
J. Dulcic and B. Grbec

MEETING REPORT

192 Highlights of the PICES Eighth Annual Meeting
Patricia Livingston

Contents

Volume 9 Number 3 September 2000

ORIGINAL ARTICLES

195 Observations on larval fish transport and retention on the Scotian Shelf in relation to geostrophic circulation
C. S. Reiss, G. Panteleev, C. T. Taggart, J. Sheng and B. deYoung

214 Distribution of minke whales in the Bellingshausen and Amundsen Seas (60°W–120°W), with special reference to environmental/physiographic variables
Fujio Kasamatsu, Paul Ensor, Gerald G. Joyce and Noritugu Kimura

224 Mid-latitude wind stress: the energy source for climatic shifts in the North Pacific Ocean
R. H. Parrish, F. B. Schwing and R. Mendelssohn

239 Ecological variations and El Niño effects off the southern coast of the Korean Peninsula during the last three decades
Suam Kim and Sukyung Kang

248 Alternating dominance of postlarval sardine and anchovy caught by coastal fishery in relation to the Kuroshio meander in the Enshu-nada Sea
Hideaki Nakata, Shigeo Funakoshi and Motohiko Nakamura

259 Behaviour of immature yellowtails (*Seriola quinqueradiata*) observed by electronic data-recording tags
Akihide Kasai, Wataru Sakamoto, Yasushi Mitsunaga and Shotaro Yamamoto

271 Computer simulations of the effects of the Sitka eddy on the migration of sockeye salmon returning to British Columbia
M. C. Healey, K. A. Thomson, P. H. Leblond, L. Huato, S. G. Hinch and C. J. Walters

RESEARCH NEWS

282 Presentation of SST images from satellite NOAA on the internet by the Fisheries Research Institute of Mie, Japan
Masahiro Kuno, Takashi Yamakawa and Hirokatsu Yamada

Contents

Volume 9 Number 4 December 2000

ORIGINAL ARTICLES

285 Pacific herring, *Clupea pallasi*, recruitment in the Bering Sea and north-east Pacific Ocean, I: relationships among different populations
Erik H. Williams and Terrance J. Quinn, II

300 Pacific herring, *Clupea pallasi*, recruitment in the Bering Sea and north-east Pacific Ocean, II: relationships to environmental variables and implications for forecasting
Erik H. Williams and Terrance J. Quinn, II

316 Relative contributions from exposed inshore and estuarine nursery grounds to the recruitment of stone flounder, *Platichythys bicoloratus*, estimated using otolith Sr:Ca ratios
Yoh Yamashita, Tsuguo Otake and Hideaki Yamada

328 A dynamic optimisation model for the behaviour of tunas at ocean fronts
David S. Kirby, Oyvind Fiksen and Paul J. B. Hart

343 The importance of retention process in upwelling areas for recruitment of *Octopus vulgaris*: the example of the Arguin bank (Mauritania)
Valerie Faure, Cheikh Abdellahi Inejih, Herve Demarcq and Philippe Cury

356 Acoustic approaches to remote species identification: a review
John K. Horne

SHORT COMMUNICATION

372 On the use of Lloyd's index of patchiness
Nicolas Bez

377 ERRATA

379 INDEXES TO VOLUME 9